arXiv:cond-mat/0307746v1 [cond-mat.str-el] 30 Jul 2003

Néel to spin-Peierls transition in the ground state of a quasi-1D Heisenberg model coupled to bond phonons

Pinaki Sengupta
Department of Physics, University of California, Davis, CA 95616
(February 2, 2008)

The spin-Peierls transition in the ground state of a quasi-one-dimensional spin-$\frac{1}{2}$ Heisenberg model coupled to adiabatic bond phonons is studied using a quantum Monte Carlo (QMC) method. The transition from a gapless Néel state to a spin-gapped Peierls state is explored in the parameter space spanned by the spatial anisotropy and the strength of spin-lattice coupling. It is found that for any finite inter-chain coupling, the transition to a dimerized Peierls ground state occurs only when the spin-lattice coupling exceeds a finite, non-zero critical value. This is in contrast to the pure 1D model, where adiabatic phonons lead to a dimerized ground state for any non-zero spin-phonon coupling. The phase diagram in the above parameter space is mapped out. No evidence is found for a region of co-existing long range magnetic order and dimerization. The vanishing of the Néel order occurs simultaneously with the setting in of the dimerization.

PACS: 75.40.Gb, 75.40.Mg, 75.10.Jm, 75.30.Ds

I. INTRODUCTION

When coupled to an elastic lattice, a spin-$\frac{1}{2}$ Heisenberg chain is unstable towards a dimerization of the lattice. The cost in elastic energy due to a distortion δ of the lattice ($\sim \delta^2$) is compensated for by the gain in the magnetic energy ($\sim \delta^{4/3}$), and the ground state is stabilized for a lattice with non-zero dimerization[1,2] and a finite spin gap. Such a transition is known as the spin-Peierls (SP) transition – in analogy with the conventional Peierls transition in a one-dimensional (1D) metal. The SP transition has been extensively studied and is believed to be fairly well understood in 1D.[1–9] In the adiabatic limit, any arbitrarily small coupling to an elastic lattice leads to a dimerized ground state. On the other hand, quantum lattice fluctuations destroy the dimerization for small spin-phonon couplings and/or large bare phonon frequencies.[6,7] The transition to a dimerized (Peierls) state occurs only when the spin-phonon coupling exceeds a finite, non-zero critical value (that depends on the bare phonon frequency). This is true for both the spin-$\frac{1}{2}$ Heisenberg and XY models with spin-phonon coupling. The discovery[10] of the quasi-1D inorganic spin-Peierls compound $\mathrm{CuGeO_3}$ has led to a resurgence in the study of spin-Peierls(SP) transition in low-dimensional spin models. The properties of $\mathrm{CuGeO_3}$ have been widely studied within the framework of a 1D spin-$\frac{1}{2}$ Heisenberg model coupled to phonons, with an additional frustrated next-nearest-neighbor interaction.[11]

Fewer studies exist for the spin-Peierls transition in two dimensions (2D) and the mechanism of the transition is not as well established as in 1D. Although the study of the Peierls transition in 2D or quasi-1D systems presents a technically much more difficult challenge, it also allows for a more realistic modeling of real materials – for $\mathrm{CuGeO_3}$, the strength of inter-chain coupling is estimated to be $J_\perp/J \approx 0.1$ experimentally[12], and its importance in any realistic modeling of $\mathrm{CuGeO_3}$ has been widely stressed.[13,14] From a purely theoretical standpoint, the study of spin-Peierls transition is interesting on its own. Unlike in 1D, the ground state of the 2D (Heisenberg) system, in the absence of spin-phonon coupling, has long range antiferromagnetic (Néel) order. It is generally believed that even for adiabatic phonons, the spin-phonon coupling has to exceed some non-zero critical value for the ground state to develop a dimerized pattern with a spin gap. What is the nature of the transition? Is there a region in the phase space where the ground state has co-existing dimerization and long range antiferromagnetic order? Furthermore, the existence of several different possible dimerization pattern in 2D means that, in principle, different dimerization patterns can be stabilized for different values of the parameters. Finally, unlike in 1D, the Peierls phase in 2D extends to finite temperatures.

The spatially isotropic 2D spin-$\frac{1}{2}$ Heisenberg model with static dimerization patterns have been studied by several authors.[15–20] By comparing the ground state energies for different patterns, these authors have tried to predict the energetically most favored dimerization pattern. Using the same strategy, the 2D tight binding model with bond-distortions[21] and the 2D Peierls-Hubbard model[22–24] have also been studied. Since in the limit of large on-site repulsion, U, the Hubbard model at half-filling reduces to the Heisenberg model, results from the Peierls-Hubbard model (in the limit of large U) are expected to be applicable to the present discussion. Unfortunately, there is no consensus among the different studies as to the nature of the dimerization pattern in the ground state. For the Peierls-Hubbard model at half-filling, Tang and Hirsch[22] find a plaquette-like distortion to be energetically favored in the limit of large U. On the other hand, Mazumdar[23] have argued that the minimum energy ground state has a "stair-like" dimerization pattern – corresponding to a wave vector $\mathbf{Q} = (\pi, \pi)$. Zhang and Prelovšek[24] agree with a dimerization pattern with $\mathbf{Q} = (\pi, \pi)$, but conclude that the ground state has dimerization only along one of the axes (staggered dimerized chains – similar to the pattern considered here). For the Heisenberg model with static dimerization, Al-Omari[19] has concluded that the ground state energy is minimized for a plaquette-like distortion of the lattice which agrees with the conclusion of Tang and Hirsch. On the other hand, Sirker *et.al.*[20] find that linear spin wave theory (LSWT) predicts a stairlike dimerization pattern to be most favored, in agreement with Mazumdar. Using LSWT, Sirker *et.al.* also find finite regions in the parameter space with co-existing long range magnetic order and non-zero dimerization. However, as pointed out by the authors, results obtained from LSWT are not reliable at large values of dimerization. The effects of inter-chain coupling was considered early on by Inagaki and Fukuyama[15,16] who studied a quasi-1D system of coupled spin-$\frac{1}{2}$ Heisenberg chains with a fixed dimerization pattern corresponding to a wave vector $\mathbf{Q} = (\pi, 0)$. By treating the inter-chain coupling in a mean-field theory, they were able map out the ground state phase diagram[15] and study the finite temperature transition[16]. Later, Katoh and Imada[17] studied in detail the nature of the transition. More recently, the effects of impurities have been studied in this model (once again with a mean-field treatment of the inter-chain coupling) which revealed a region of co-existing Peierls and antiferromagnetic orders.[25–30] In addition to this, the quasi-1D XY model with a $\mathbf{Q} = (\pi, \pi)$ dimerization pattern has recently been studied[31,32] by an extension of the Jordan-Wigner transformation in 2D.[33–35] The effects of quantum phonons in the isotropic 2D model have also been studied,[36] where the authors find that there is no evidence of a transition to the Peierls state for a wide range of values of the bare phonon frequency and the spin-phonon coupling. This is consistent with previous finding[37] for the same model that the spin wave spectrum along the Brillouin zone boundary is qualitatively similar to that for the pure Heisenberg model.

The present work aims to investigate the SP transition in a spin-$\frac{1}{2}$ quasi-1D Heisenberg antiferromagnet coupled to static $\mathbf{Q} = (\pi, \pi)$ bond phonons with varying inter-chain coupling. The elastic energy due to the lattice distortion is considered as well as the change in magnetic energy. The nature of the ground state is explored for different values of the inter-chain coupling and the elastic constant associated with the bond distortions and the ground state phase diagram in this phase space is mapped out.

The rest of the paper is organized as follows. In Section II, the model Hamiltonian and the Stochastic Series Expansion (SSE) QMC method used to study it are introduced. The results of the simulations are presented in Section III. Section IV concludes with a summary of the results.



FIG. 1. The spin part of the ground state energy as a function of the system size for four values of δ, and α=0.25.

II. MODEL AND SIMULATION TECHNIQUES

A quantum Monte Carlo (QMC) method has been used to study a quasi-1D Heisenberg model with spin-phonon coupling. The model is given by the Hamiltonian

$$\begin{aligned} H = J \sum_{i,j} (1 + \lambda u_{i,j}) \mathbf{S}_{i,j} \cdot \mathbf{S}_{i+1,j} + \frac{1}{2} K \sum_{i,j} u_{i,j}^2 \\ + J_\perp \sum_{i,j} \mathbf{S}_{i,j} \cdot \mathbf{S}_{i,j+1}, \end{aligned} \tag{1}$$

where $J_\perp$ is the inter-chain coupling, λ is the strength of the spin-phonon coupling (restricted to be only along the chains), $u_{i,j}$'s are the distortions of the bond lengths and K is the elastic constant for the distortions. Following the experimentally observed[38] dimerization pattern in CuGeO$_3$, the bond length distortions are chosen to be of the form

$$u_{i,j} = (-1)^{i+j} \delta$$

This amounts to choosing a fixed dimerization pattern corresponding to the wave-vector $\mathbf{Q} = (\pi, \pi)$. As in previous works, the bond distortions can be rescaled by the spin-phonon coupling strength, λ. This reduces the Hamiltonian to



FIG. 2. The spin part and the total ground state energy as a function of the bond distortion for 3 different values of the elastic constant, ζ, and fixed α=0.25. The system size is N=64x16.

$$H = \sum_{i,j}(1+(-1)^{i+j}\delta)\mathbf{S}_{i,j}\cdot\mathbf{S}_{i+1,j} + N\delta^2/2\zeta$$
$$+\alpha\sum_{i,j}\mathbf{S}_{i,j}\cdot\mathbf{S}_{i,j+1}, \qquad (2)$$

where $\zeta = \frac{\lambda^2 J}{K}$, $\alpha = J_\perp/J$ and N is the size of the lattice. The static approximation for the displacements makes the computational task easier. The following strategy is adopted. The simulations are carried out for the spin variables for several different $\{\alpha, \delta\}$ parameter sets. This produces the spin energy of the system as a function of δ for a fixed α. Next the elastic energy with a particular ζ is added and the total energy is minimized to obtain the value of the ground state distortion for the given set of parameters $\{\alpha, \zeta\}$. This is repeated for different sets of $\{\alpha, \zeta\}$ to obtain the ground state phase diagram in the parameter space spanned by α and ζ.

The stochastic series expansion (SSE) QMC method has been used to study the present model. The SSE method[39−41] is a finite-temperature quantum Monte Carlo method based on importance sampling of the diagonal elements of the Taylor expansion of $e^{-\beta H}$, where β is the inverse temperature $\beta = J/T$. Ground state expectation values can be obtained using sufficiently large values of β, and there are no approximations beyond statistical errors. Using the recently developed "operator loop update",[41] it is possible to explore the phase space of the Hamiltonian (2) in an efficient manner.

III. RESULTS



FIG. 3. The equilibrium ground state bond distortion as a function of the spin-lattice coupling for different values of the inter-chain coupling.

For a system of weakly coupled Heisenberg chains, it was shown recently that the estimates for various observables for a spatially anisotropic system depend non-monotonically on the system size for square ($L_x = L_y$) geometry.[42] One has to go to rectangular ($L_x \neq L_y$) geometries to obtain monotonic behavior of the numerical results for extrapolating to the thermodynamic limit. Similar effects are expected in the present model for $\alpha \ll 1$. Hence rectangular lattices with the aspect ratio $L_x = 4L_y$have been studied, with $L_x = 16-64$. An inverse temperature of $\beta = 8L_x$ was found to be sufficient for the observables to have converged to their ground state values. Nine different values of the inter-chain coupling were considered, $\alpha = 0.01, \ldots, 1.00$.

Fig.1 shows the finite-size dependence of the QMC data for the spin part of the ground state energy for four different values of the bond length distortion, δ. The values of δ chosen correspond to a Néel ground state (δ=0.04, 0.10), a Peierls ground state (δ=0.20), and a ground state close to the critical δ for Néel to spin-Peierls transition in the pure spin system (the phase transition in the spin part of the Hamiltonian (2) is discussed at the end of the current section). The leading order finite-size correction to the ground state energy is seen to be $\sim 1/L_x^3$ for small values of δ corresponding to a Néel-ordered ground state. This is similar to that observed for the pure 2D Heisenberg model.[40] On the other hand, for a spin-gapped ground state (δ=0.20), the energy scales exponentially with system-size. Close to the critical point, extrapolation to the thermodynamic limit becomes difficult due to cross-over effects. Instead, the data from the largest system size studied have been used to map out the phase diagram. Fortunately, the data for the largest system sizes studied are found to be well converged – the fractional difference in the energy for the two largest system sizes studied is $\sim 10^{-5}$. This observed convergence allows for reliable estimation of ground state properties in the thermodynamic limit based on the data from the largest system size – any finite-size effects on such estimates are expected to be small.

The strategy implemented to extract the ground state bond distortion is qualitatively demonstrated in fig.2. The total ground state energy is obtained by adding the elastic energy contribution to the spin part of the energy obtained from the simulations. The plot shows the spin part of the energy, as well as the total ground state energy as a function of the bond length distortion, δ, for three different values of the elastic energy constant, ζ, and a fixed value of the inter-chain coupling (α=0.25). For large ζ, the total energy is minimized for a non-zero value of the bond length distortion, δ, whereas for small ζ, a uniform ground state with δ=0 is energetically favored. The behavior of the total energy near the critical ζ is also shown. The ground state distortion in bond length is obtained by numerically differentiating the total energy data and solving for $\frac{\partial \mathrm{E}^{tot}}{\partial \delta}|_{\delta^{gs}} = 0$. In principle, one can also fit a polynomial to the QMC data to get $\mathrm{E}^{spin}(\delta)$ and add to it the elastic energy term to get $\mathrm{E}^{tot}(\delta)$. Then the ground state distortion can be obtained as a continuous function of δ by solving $\frac{\partial \mathrm{E}^{tot}}{\partial \delta}|_{\delta^{gs}} = 0$. However, in practice, the earlier method is found to be more reliable because of the uncertainty in the order of the polynomial fit.

Fig.3 shows the equilibrium distortion in the bond lengths in the ground state of the system as a function of the elastic energy parameter, ζ, at fixed values of α, obtained as described above. For small values of ζ, the tendency towards dimerization is suppressed, and a uniform (Néel ordered) ground state with uniform bond lengths is stabilized. As ζ is increased above a critical value, ζ_c, there is a transition to a ground state with a finite, non-zero dimerization. For $\zeta > \zeta_c$, the equilibrium ground state distortion increases monotonically with ζ. The critical value ζ_c required for a transition to a dimerized ground state depends on the value of the inter-chain coupling, α and increases with increasing α. The data indicate that the transition is a discontinuous (first order) quantum phase transition for any finite value of inter-chain coupling.



FIG. 4. The ground-state phase diagram in the parameter space of the inter-chain coupling and the spin-lattice coupling strength.

The results from fig.3 are summarized in fig.4, which shows the phase diagram for the system in the phase space spanned by the parameters ζ and α. For small ζ and/or large α, the ground state of the system is Néel ordered with zero bond distortion, while for large ζ and/or small α, the ground state is dimerized with a finite spin gap. The phase boundary is roughly linear over the range of values of α studied. However, it is well known that in the 1D limit ($\alpha = 0$), the critical elastic constant should vanish ($\zeta_c = 0$). Unfortunately, the numerical data becomes too noisy in this limit for an accurate estimate of the critical coupling. For the quasi-1D XY model, it was shown that ζ_c goes as $-1/\ln\alpha$ for $\alpha \to 0$. On the other hand, results for a system of coupled (undimerized) Heisenberg chains show that the magnetization vanishes as a power law in α with a logarithmic correction.[42] A similar behavior is expected for the critical coupling in the present model.

This brings up the interesting possibility of having a region in the (ζ, α) parameter space where the ground state has co-existing Néel order and non-zero dimerization. Such co-existence has been shown to exist in the presence of doping.[25–30] For this purpose, the static spin susceptibility defined as

$$\chi(\mathbf{q}) = \frac{1}{N} \sum_{\langle i,j \rangle} e^{i\mathbf{q}\cdot(\mathbf{r}_j - \mathbf{r}_j)} \int_0^\beta d\tau \langle S_j^z(\tau) S_i^z(0) \rangle. \tag{3}$$

have been studied for the spin part of the Hamiltonian (2)(without the elastic energy term).

$$\begin{aligned} H = \sum_{i,j} (1 + (-1)^{i+j}\delta) \mathbf{S}_{i,j} \cdot \mathbf{S}_{i+1,j} + \\ + \alpha \sum_{i,j} \mathbf{S}_{i,j} \cdot \mathbf{S}_{i,j+1}, \end{aligned} \tag{4}$$

If the ground state has long range antiferromagnetic order, the staggered ($\mathbf{Q} = (\pi, \pi)$) susceptibility scaled by the system size ($\chi(\pi, \pi)/N$), for a finite system will increase with increasing system size, diverging in the thermodynamic limit. On the other hand, if the ground state has a finite spin gap, $\chi(\pi, \pi)/N$ will vanish in the limit of infinite system size. This qualitative criterion can be expressed in a more quantitative manner by noting that the ground state of the above Hamiltonian undergoes a continuous transition from a Néel ordered state with long range antiferromagnetic order to a spin-gapped, dimerized phase with no long range magnetic order as δ is increased beyond a finite, non-zero critical value, δ^*, that depends on the inter-chain coupling α. The transition belongs to the universality class of the 3D Heisenberg model[43]. Finite-size scaling[44] predicts that for such a transition, the finite-size susceptibility at the critical δ scales with the system size as

$$\chi(L_x) \sim L_x^{2-\eta},$$

for a rectangular lattice of dimension $N = L_x \text{x} L_y$. This implies that on a plot of $\chi(\pi,\pi)/L_x^{2-\eta}$, the curves for different system sizes will intersect at the critical δ. The value of the critical exponent η is known to a high degree of accuracy ($\eta \approx 0.037$).[45] Fig. 5 shows $\chi(\pi,\pi)/L_x^{2-\eta}$ as a function of δ for a fixed value of $\alpha = 0.25$ for several different system sizes. For small δ, the scaled susceptibility increases with increasing system size, indicating the presence of long range magnetic order. For larger values of δ, the scaled susceptibility goes to zero with increasing system size, signaling the opening up of a spin gap. From the data, the critical value of δ is estimated to be $\delta^* \approx 0.16$. This value of the bond length distortion is less than the jump in δ at the transition point, ζ_c. This means that for $\zeta > \zeta_c$, the ground state is dimerized with $\delta^{gs} > \delta^*$ and has no long range magnetic order. On the other hand, for $\zeta < \zeta_c$, the a uniform ground state is energetically favored that has zero dimerization ($\delta^{gs} = 0$), and long range magnetic order (the Néel state). For no values of ζ is a ground state with $0 < \delta^{gs} < \delta^*$ ever favored energetically. Hence the transition to the dimerized phase is accompanied by the disappearance of magnetic order and there is no region of co-existing dimerization and magnetic order. This is true for the present model. Whether it is possible to have ground states with co-existing magnetic order and dimerization in other models (with different dimerization patterns) remains to be seen.



FIG. 5. The ground-state static staggered magnetic susceptibility as a function of bond distortion for a fixed value of the inter-chain coupling, $\alpha = 0.25$.

IV. SUMMARY

A quantum Monte Carlo method has been used to study a quasi-1D Heisenberg model coupled to static bond phonons. Motivated by experimental observations in the inorganic quasi-1D spin-Peierls compound CuGeO$_3$, the bond distortions are restricted to be only along the chains, and a single dimerization pattern, corresponding to the wave vector $\mathbf{Q} = (\pi,\pi)$, is considered. It is found that unlike the pure 1D case, for any non-zero inter-chain coupling α, the transition to a dimerized spin-Peierls ground state occurs only when the spin-lattice coupling strength, ζ, exceeds a finite, non-zero critical value, ζ_c, even for the static phonons considered here. For $\zeta < \zeta_c$, the ground state has long range Néel order and zero spin gap, whereas for $\zeta > \zeta_c$, the ground state develops a finite dimerization accompanied by the opening up of a spin gap. The transition is found to be a discontinuous (first order) quantum phase transition. The value of the critical coupling depends on the strength of the inter-chain coupling, increasing monotonically with α. The phase diagram in the parameter space of α and ζ is mapped out. Finally it is found that in the present model, the transition to the dimerized Peierls state is accompanied by the vanishing of magnetic order, and that there is no region of co-existing magnetic order and non-zero dimerization.

V. ACKNOWLEDGMENTS

The author would like to thank Anders Sandvik and Rajiv Singh for useful discussions. This work was supported in part by NSF grant number DMR-9986948. Simulations were carried out at the IBM SP facility at NERSC, Berkeley.

---

[1] E. Pytte, Phys. Rev. B **10**, 4637 (1974).

[2] M. C. Cross, and D. S. Fisher, Phys. Rev. B **19**, 402 (1979).

[3] G. S. Uhrig, Phys. Rev. B **57**, R14004 (1998).

[4] C. Gros and R. Werner, Phys. Rev. B **58**, R14677 (1998).

[5] G. Wellein, H. Fehske, and A. P. Kampf, Phys. Rev. Lett. **81**, 3956 (1998).

[6] A. W. Sandvik and D. K. Campbell, Phys. Rev. Lett. **83**, 195 (1999).

[7] R. J. Bursill, R. H. McKenzie and C. J. Hamer, Phys. Rev. Lett. **83**, 408 (1999).

[8] A. Weiße, G. Wellein, and H. Fehske, Phys. Rev. B **60**, 6566 (1999).

[9] C. Raas, U. Löw, G. S. Uhrig, and R. W. Kühne, Phys. Rev. B **65**, 144438 (2002).

[10] M. Hase, I. Terasaki, and K. Uchinokura, Phys. Rev. Lett. **70**, 3651 (1993).

[11] J. Riera, and A. Dobry, Phys. Rev. B **51**, 16098 (1995).
[12] M. Nishi, O. Fujita, and J. Akimitsu, Phys. Rev. B **50**, 6508 (1994).
[13] G. S. Uhrig, Phys. Rev. Lett. **79**, 163 (1997).
[14] G. Bouzerar, Ö. Legeza, and T. Ziman, Phys. Rev. B, **60**, 15278 (1999).
[15] S. Inagaki, and H. Fukuyama, J. Phys. Soc. Jpn. **52**(10), 3620 (1983).
[16] S. Inagaki, and H. Fukuyama, J. Phys. Soc. Jpn. **57**(2), 1435 (1988).
[17] N. Katoh, and M. Imada, J. Phys. Soc. Jpn. **62**, 3728 (1993); *ibid.* **63**, 4529 (1994).
[18] A. Koga, S. Kumada, and N. Kawakami, J. Phys. Soc. Jpn. **68**, 642 (1998); *ibid.* **68**, 2373 (1999).
[19] A. Al-Omari, J. Phys. Soc. Jpn. **69**(10), 3387 (2000).
[20] J. Sirker, A. Klümper, and K. Hamacher, Phys. Rev. B **65**, 134409 (2002).
[21] Y. Ono, and T. Hamano, J. Phys. Soc. Jpn. **69**(6), 1769 (2000).
[22] S. Tang, and J. E. Hirsch, Phys. Rev. B, **37**, 9546 (1988).
[23] S. Mazumdar, Phys. Rev. B, **36**, 7190 (1987).
[24] F. C. Zhang, and P. Prelovšek, Phys. Rev. B, **37**, 1569 (1988).
[25] M. Mostovoy, D. Khomskii, and J. Knoester, Phys. Rev. B **58**, 8190 (1998).
[26] M. Saito, J. Phys. Soc. Jpn. **68**(9), 2898 (1999).
[27] E. Sørensen, and I Affleck, D. Augier, and D. Poilblanc, Phys. Rev. B **58**, R14701 (1998).
[28] A. Dobry, P. Hansen, J. Riera, D. Augier, and D. Poilblanc, Phys. Rev. B **60**, 4065 (1999).
[29] M. Fabrizio, R. Mélin, and J. Souletie, Eur. Phys. J. B **10**, 607 (1999).
[30] R. Mélin, Eur. Phys. J. B **16**, 261 (2000); *ibid.* **18**, 263 (2000).
[31] Y. Ji, J. Qi, J.-X. Li, and C.-D. Gong, J. Phys: Cond. Mat. **9**, 2259 (1997).
[32] Q. Yuan, Y. Zhang, and H. Chen, Phys. Rev. B **64**, 12414 (2001).
[33] Y. R. Wang, Phys. Rev. B, **43**, 3786 (1991).
[34] Y. R. Wang, Phys. Rev. B, **46**, 151 (1992).
[35] M. Azzouz, and C. Bourbonnais, Phys. Rev. B **53**, 5090 (1996).
[36] C. H. Aits, and U. Löw, cond-mat/0306745.
[37] P. Sengupta, R. T. Scalettar, and R. R. P. Singh, Phys Rev. B **66**, 144420 (2002).
[38] K. Hirota, D. E. Cox, J. E. Lorenzo, G. Shirane, J. M. Tranquada, M. Hase, K. Uchinokura, H. Kojima, Y. Shibuya, and I. Tanaka, Phys. Rev. Lett. **73**, 736 (1994).
[39] A. W. Sandvik, J. Phys. A **25**, 3667 (1992).
[40] A. W. Sandvik, Phys. Rev. B **56**, 11678 (1997).
[41] A. W. Sandvik, Phys. Rev. B **59**, 14157 (1999).
[42] A. W. Sandvik, Phys. Rev. Lett. **83**, 3069 (1999).
[43] S. Chakravarty, B. I. Halperin and D. R. Nelson, Phys. Rev. Lett. **60**, 1057 (1988); Phys. Rev. B **39**, 2344 (1989).
[44] For a review of finite-size scaling, see: M. N. Barber in *Phase Transitions and Critical Phenomena, Vol. 8*, ed. Domb and Lebowitz (Academic Press, 1983).
[45] M. Campostrini, M. Hasenbusch, A. Pelissetto, P. Rossi, and E. Vicari, Phys. Rev. B **65**, 144520 (2002).